



FOR RESIDENTIAL USE

Net Wt. 50 lbs.
Treats Up To 14,000 Sq. Ft.

REPAR BROAD SPECTRUM WEED & FEED

- For use on warm and cool season turf types as listed.
- Controls dandelions, clover, dollarweed, chickweed and 200 common lawn weeds (as listed).
- For a thick green weed free lawn.

ACTIVE INGREDIENTS

Chlorflurenol-Methyl	0.17%
Dicamba.....	0.13%

OTHER INGREDIENTS.....

.....	99.70%
Total	100.00%

Fertilizer and Carrier information will be supplied by registrants/sub-registrants to States and other regulatory authorities who have jurisdiction over these ingredients.

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

Causes moderate eye irritation. Avoid contact with eyes, skin or clothing.

Wear goggles, face shield or safety glasses and waterproof gloves. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco, or using toilet.

FIRST AID

If In Eyes	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing. Call a poison control center or doctor for treatment advice.
If on Skin or Clothing	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15 - 20 minutes. Call a poison control center or doctor for treatment advice.
If Swallowed	Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to by a poison control center or doctor.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

HOT LINE NUMBER

For 24-Hour Medical Emergency Assistance (Human or Animal) or Chemical Emergency Assistance (Spill, Leak, Fire or Accident) call **CHEMTREC** at 1-800-424-9300.

Manufactured for:

Repar Corporation

P.O. Box 4321 • Silver Spring, MD 20914

EPA Reg. No. 69361-3

EPA Est. No. 51274-1

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area. Sweeping any product that lands on a driveway, sidewalk, or street, back onto the treated area of the lawn or garden will help to prevent run off to water bodies or drainage systems.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact any person or pet, either directly or through drift. Keep people and pets out of the area during application. Do not allow people or pets to enter the treated area until dusts have settled.

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass. Not for use on turf being grown for sale or other commercial use as sod, or for commercial seed production, or for research purpose.

WEED CONTROL

Repar Broad Spectrum Weed and Feed controls a wide range of perennial and annual weeds including Dandelion, Buckhorn Plaintain, Common Chickweed, White and Red Clover, Hop Clover, Marestalk, Heal All, Henbit, Ground Ivy, Knotweed, Black Medic, Purslane, Yellow Rocket, Thistles, & other listed Broadleaf Weeds.

WEED LIST

Annuals

Amaranth, Spiny (Spiny Pigweed)	Goosefoot, Nettleleaf	Pennycress, Field (Fanweed,	Senna, Coffee
Aster, Slender	Henbit	Frenchweed,	Sesbania, Hemp
Bedstraw	Jimsonweed	Stinkweed)	Shephardspurse
Beggarweed, Florida	Knotweed	Pepperweed, Virginia (Peppergrass)	Sicklepod
Broomweed, Common	Kochia (triazine resistant)	Pigweed, Prostrate	Sida, Prickly (Teaweed)
Buckwheat, Wild	Kochia	Pigweed, Redroot (Carelessweed)	Smartweed, Green
Buffalobur	Ladysthumb	Pigweed, Rough (Carelessweed)	Smartweed, Pennsylvania
Burclover, California	Lambsquarters, Common	Pigweed, Rough (Carelessweed)	Sneezeweed, Bitter
Burcucumber	Lambsquarters, (triazine resistant)	Pigweed, Rough (triazine resistant)	Sowthistle, Annual
Buttercup, Roughseed	Lettuce, Prickly	Pigweed, Tumble	Sowthistle, Spiny
Carpetweed	Mallow, Common	Poorjoe	Spanishneedles
Catchfly, Nightflowering	Mallow, Venice	Puncturevine	Spikeweed, Common
Chamomile, Corn	Marestalk	Purslane, Common	Spurge
Chickweed, Common	(Horseweed)	Pusley, Florida	Spurr, Corn
Clovers, Annual	Mayweed	Radish, Wild	Starbur, Bristly
Cockle, Corn	Medic, Black	Ragweed, Common	Stumpweed, Rough
Cocklebur, Common	Morningglory, Ivyleaf	Ragweed, Giant (Buffalowed)	Sunflower, Common (Wild)
Croton, Tropic	Morninggorty, Tall	Ragweed, Lance-Leaf	Sunflower, Volunteer
Croton, Woolly	Mustard, Tansy	Rubberweed, Bitter (Bitterweed)	Thistle, Russian
Daisy, English	Mustard, Wild		Thistle, Milk
Eveningprimrose,	Mustard (Yellowtops)		Thistle, Musk
Cutleaf	Nightshade, Black		Thistle, Plumeless
Fleabane, Annual			

Biennials

Burdock, Common	Geranium	Plaintain, Bracted	Thistle, Bull
Carrot, Wild (Queen Anne's Lace)	Gromwell	Ragwort, Tansy	Thistle, Milk
Cockle, White	Knapweed, Diffuse	Starthistle, Yellow	Thistle, Musk
Eveningprimrose, Common	Knapweed, Spotted	Sweetclover	Thistle, Plumeless
	Mallow, Dwarf	Teasel	

Perennials

Alfalfa*	Clover, Hop*	Milkweed, Common	Sowthistle, Perennial
Artichoke, Jerusalem	Dandelion, Common*	Milkweed, Honeyvine	Spurge, Leafy
Aster, Spiny	Dock, Broadleaf* (Bitterdock)	Milkweed, Western Whorled	Sundrop, Halfshrub (Eveningprimrose)
Aster, Whiteheath	Dock, Curly*	Nettle, Stinging	Thistle, Canada
Bedstraw, Smooth	Dogbane, Hemp	Nightshade, Silverleaf	Toadflax Dalmation
Bindweed, Field	Dogfennel*	(White Horsenettle)	Tropical Soda Apple
Bindweed, Hedge	(Cypressweed)	Onion, Wild	Trumpetcreeper (Buckvine)
Bindweed, Texas	Fern, Bracken	Pennywort (Dollar- weed)	Vetch
Bursage*	Garlic, Wild	Plantain, Broadleaf*	Violet, Wild
(Bur Ragweed, Lakeweed, Povertyweed)	Goldenrod, Canada	Plantain, Buckhorn	Waterhemlock
Bursage, Woolyleaf (Lakeweed)	Goldenrod, Missouri	Pokeweed	Waterprimrose Creeping
Buttercup, Tall	Goldenweed, Common	Ragweed, Western	Woodsorrel, Creeping* (Common Yellow)
Campion, Bladder	Hawkweed	Sericia Lespedeza	Wormwood, Common
Chickweed, Field	Henbane, Black	Redvine	Wormwood, Louisiana
Chickweed, Mouseear (Canada)	Horsenettle, Carolina	Smartweed, Swamp	Yankeeweed*
Chicory	Ironweed	Snakeweed, Broom	Yarrow, Common
	Ivy, Ground	Sorrel, Red* (Sheep Sorrel)	
	Knapweed, Black	Sowthistle	
	Knapweed, Russian		
	Milkweed, Climbing		

* Noted perennials may be controlled using Repar Broad Spectrum Weed and Feed at rates lower than those directed for other listed perennial weeds.

Woody

Ailanthus	Ivy, Poison	Kudzu	Sagebrush, Fringe
Sassafras	Gallberry	Serviceberry	Wild Grape
Oak, Poison	Sumac	Brazilian Pepper	Tarbrush
Honeysuckle	Hornbeam	Witchhazel	Huckleberry
Huisache			

For Use on Established Turfs, including sensitive species such as Centipede and St. Augustine, except Desirable Clovers, Bent, Dichondra and Carpet Grass.

EFFECTIVE APPLICATION OF REPAR BROAD SPECTRUM WEED AND FEED FOR MOST WEED LEAVES SHOULD BE DAMP FROM RAIN, DEW OR SPRINKLING TO ALLOW THE PRODUCT TO STICK TO THE LEAF. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL.

WHAT TO LOOK FOR:

Repar Broad Spectrum Weed and Feed, properly applied, first reduces or eliminates the formation of most flower heads and the seeds they produce and then attacks and removes the mature weed itself. Look for an absence or stunting of seed heads in flowering weeds such as clover or dandelions. Some weeds may also show twisting stems and dwarfed appearance with curling of the leaves.

Repar Broad Spectrum Weed and Feed is designed with an excellent turf nutrient diet to allow grass plants to gradually fill in the areas around dying weeds. This allows for a more natural recovery of the turf and minimizes reinfestations of bare areas by wind-borne weed seeds. Applied as directed, the rich green color will not be affected.

UNDER NORMAL GROWING CONDITIONS, WEED KILL SHOULD BE COMPLETE WITHIN 21 TO 35 DAYS

SPREADER SETTINGS

Spreader	Rate: 3.5#/M	Pattern Setting	Width
Earthway Broadcast	11	-	8
Lely	4 1/2	III - B	42
Lesco Broadcast	4 1/2	0.0	11
Prizelawn Broadcast	4 1/4	A	7
Scotts Broadcast	1	5	10
Spyker	3	-	9
Vicon	29	-	4
Cyclone Broadcast	3 1/4	-	5

Important: These are approximate settings. Age and condition of your spreader can cause wide variation. Be sure to calibrate your spreader with each application.

WHEN TO APPLY:

Apply any time weeds are actively growing and temperature is above 60°F. Do not apply when turf is under stress from heat, excessive moisture or drought. Control is quicker and more complete if applied in Spring prior to weed flower formation. Control is slower if applied when weather is cool. On newly seeded areas, wait until grass has been cut two or three times before using. Delay planting new grass for at least three weeks as the active ingredients may temporarily inhibit seed germination.

HOW TO APPLY:

EFFECTIVE APPLICATION OF REPAR BROAD SPECTRUM WEED AND FEED REQUIRES THE WEED LEAVES TO BE DAMP FROM RAIN, DEW OR SPRINKLING PRIOR TO APPLICATION. A LACK OF MOISTURE ON THE WEED LEAVES WILL REDUCE WEED CONTROL. Apply evenly over the entire lawn at the rate of 3.5 lbs./1,000 sq. ft. Rain or heavy sprinkling with 24 hours of application will temporarily delay weed control. Mowing within 24 hours may reduce effectiveness. Wait at least 45 days before making a second application. Do not make more than two applications per year.

Use the suggested spreader settings as a guide being sure to check your spreader at the beginning of the application. Adjust your spreader if necessary. Be sure to shut off spreader while turning or stopping to prevent accidental over application and burning of grass. **SPECIAL NOTE:** Repar Broad Spectrum Weed And Feed is designed to be absorbed by plant leaves. Accidental application of this product to the foliage of ornamentals such as roses may result in the loss of flowers. For this reason, extra care should be taken when using a broadcast or rotary type spreader.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in original container in a locked storage area. Keep container closed and away from food, feedstuffs and domestic water supplies.

PESTICIDE DISPOSAL AND CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container. If empty: Place in trash or offer for recycling if available. If partly filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

IMPORTANT: READ BEFORE USE:

Read the entire Directions for Use and the Warranty Disclaimer and Limitation of Liability before using this product. If terms are not acceptable, return the unopened product container at once. By using this product, User or Buyer accepts the following Warranty Disclaimer and Limitation of Liability:

WARRANTY DISCLAIMER and LIMITATION of LIABILITY

Manufacturer warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated on the label when used in strict accordance with the directions. To the extent consistent with applicable law manufacturer makes NO OTHER EXPRESS OR IMPLIED WARRANTY OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. To the extent consistent with applicable law, it is Manufacturer's intent to LIMIT ANY LIABILITY FOR SPECIAL, CONSEQUENTIAL OR INCIDENTAL ECONOMIC DAMAGES to refund of purchase price or replacement of product, at Buyer's choice. To the extent consistent with applicable law, Manufacturer DISCLAIMS ANY LIABILITY FOR COMPENSATORY OR OTHER DAMAGES ARISING OUT OF ANY USE CONTRARY TO LABEL DIRECTIONS. Use contrary to label directions is not permitted.

LEGAL RIGHTS:

THIS LIMITED WARRANTY GIVES YOU SPECIFIC LEGAL RIGHTS: YOU MAY HAVE OTHER RIGHTS THAT VARY FROM STATE JURISDICTION TO STATE JURISDICTION.

Manufactured by:

Nelson Plant Food
P.O. Box 638
Bellville, TX 77418